

Tamarisk (*Tamarix* spp.) Biology, Control, and Impacts Reference List

Bailey JK, Schweitzer JA, Whitham T. (2001) Salt Cedar Negatively Affects Biodiversity of Aquatic Macroinvertebrates. *Wetlands* 21(3):442-447.

Barz D, Watson RP, Kanney JF, Roberts JD, Groenveld DP. (2009) Cost/Benefit Considerations for Recent Saltcedar Control, Middle Pecos River, New Mexico. *Environmental Management* 43:282-298.

Bateman HL, Chung-MacCoubrey A, Snell HL. (2008) Impact of Non-Native Plant Removal on Lizards in Riparian Habitats in the Southwestern United States. *Restoration Ecology* 16(1):180–190.

Bay RF, Sher AA. (2008) Success of Active Revegetation after *Tamarix* Removal in Riparian Ecosystems of the Southwestern United States: A Quantitative Assessment of Past Restoration Projects. *Restoration Ecology* 16(1):113-128.

Bean DW, Dudley T, Keller JC. (2007) Seasonal Timing of Diapause Induction Limits the Effective Range of *Diorhabda elongata deserticola* (Coleoptera: Chrysomelidae) as a Biological Control Agent for Tamarisk (*Tamarix* spp.). *Environmental Entomology* 31(1):15-25.

Beauchamp VB, Stromberg JC. (2007) Flow Regulation of the Verde River, Arizona Encourages *Tamarix* Recruitment but has Minimal Effect on *Populus* and *Salix* Stand Density. *Wetlands* 27(2):381-389.

Beauchamp VB, Stromberg JC, Stutz JC. (2005) Interactions between *Tamarix ramosissima* (saltcedar), *Populus fremontii* (cottonwood), and mycorrhizal fungi: Effects on seedling growth and plant species coexistence. *Plant and Soil* 275:221-231.

Birken AS, Cooper DJ. (2006) Processes of *Tamarix* Invasion and Floodplain Development Along the Lower Green River, Utah. *Ecological Applications* 16(3):1103-1120.

Brand LA, White GC, Noon BR. (2008) Factors Influencing Species Richness and Community Composition of Breeding Birds in a Desert Riparian Corridor. *The Condor* 110 (2):199-210.

Brotherson JD, Field D. (1987) *Tamarix*: Impacts of a Successful Weed. *Rangelands* 9(3):110-112.

Dennison PE, Nagler PL, Hultine KR, Glenn EP, Ehleringer JR. (2009) Remote monitoring of tamarisk defoliation and evapotranspiration following saltcedar leaf beetle attack. *Remote Sensing of Environment* 113:1462-1472.

Dewine JM, Cooper DJ. (2008) Canopy shade and the successional replacement of tamarisk by native box elder. *Journal of Applied Ecology* 45:505–514.

DeWine JM, Cooper DJ. (2009) Habitat Overlap and Facilitation in Tamarisk and Box Elder Stands: Implications for Tamarisk Control Using Native Plants. *Restoration Ecology* 1-10.

- Durst SL, Theimer TC, Paxton EH, Sogge MK. (2008) Temporal variation in the arthropod community of desert riparian habitats with varying amounts of saltcedar (*Tamarix ramosissima*). *Journal of Arid Environments* 72:1644–1653.
- Durst SL, Theimer TC, Paxton EH, Sogge MK. (2008) Age, Habitat, and Yearly Variation in the Diet of a Generalist Insectivore, the Southwestern Willow Flycatcher. *The Condor* 110(3): 514–525.
- Evangelista P, Kumar S, Stohlgren TJ, Crall AW, Newman GJ. (2007) Modeling Aboveground Biomass of *Tamarix ramosissima* in the Arkansas River Basin of Southeastern Colorado, USA. *Western North American Naturalist* 67(4): 503–509.
- Field Guide for Managing Saltcedar. (2010) USDA Forest Service, Southwestern Region. TP-R3-16-2. 12pp.
- Finch DM, Yong W. (2000) Landbird Migration in Riparian Habitats of the Middle Rio Grande: A Case Study. *Studies in Avian Biology* 20:88–98.
- Gaskin JF, Kazmer DJ. (2009) Introgression between invasive saltcedars (*Tamarix chinensis* and *T. ramosissima*) in the USA. *Biological Invasions* 11:1121–1130.
- Glenn EP, Nagler PL. (2005) Comparative Ecophysiology of *Tamarix ramosissima* and Native Trees in Western U.S. Riparian Zones. *Journal of Arid Environments* 61:419–446
- Going BM, Dudley TL. (2008) Invasive riparian plant litter alters aquatic insect growth. *Biological Invasions* 10:1041–1051.
- Hamada Y, Stow DA, Coulter LL, Jafolla JC, Hendricks LW. (2007) Detecting Tamarisk species (*Tamarix* spp.) in riparian habitats of Southern California using high spatial resolution hyperspectral imagery. *Remote Sensing of Environment* 109:237–248.
- Harms RS, Hiebert RD. (2006) Vegetation Response Following Invasive Tamarisk (*Tamarix* spp.) Removal and Implications for Riparian Restoration. *Restoration Ecology* 14(3):461–472
- Hayes II WE, Walker LR, Powell EA. (2009) Competitive abilities of *Tamarix aphylla* in southern Nevada. *Plant Ecology* 202:159–167.
- Hudgeons JL, Knutson AE, Heinz KM, DeLoach CJ, Dudley TL, Pattison RR, Kiniry JR. (2007) Defoliation by Introduced *Diorhabda elongata* leaf beetles (Coleoptera: Chrysomelidae) Reduces Carbohydrate Reserves and Regrowth of *Tamarix* (Tamaricaceae). *Biological Control* 43:213–221.
- Kennedy TA, Finlay JC, Hobbie SE. (2005) Eradication of Invasive *Tamarix ramosissima* Along a Desert Stream Increases Native Fish Density. *Ecological Applications* 15(6):2072–2083.
- Kimball BA, Perry KR. (2008) Manipulating Beaver (*Castor canadensis*) Feeding Responses to Invasive Tamarisk (*Tamarix* spp.). *Journal of Chemical Ecology* 34:1050–1056.
- Lesica P, Miles S. (2001) Tamarisk Growth at the Northern Margin of Its Naturalized Range in Montana, USA. *Wetlands* 21(2):240–246.

McDaniel KC, DiTomaso JM, Duncan CA. (2005). Tamarisk or saltcedar, *Tamarix* spp. JK Clark and CL Duncan, eds. Pages 198–222 in, Assessing the Economic, Environmental and Societal Losses from Invasive Plants on Rangeland and Wildlands. Weed Science Society of America.

Milbrath LR, Deloach CJ, Jack C, Tracy JL. (2007) Overwintering Survival, Phenology, Voltinism, and Reproduction Among Different Populations of the Leaf Beetle *Diorhabda elongata* (Coleoptera: Chrysomelidae). *Environmental Entomology*, 36(6):1356-1364.

Moline AB, Poff NL. (2008) Growth of an invertebrate shredder on native (*Populus*) and non-native (*Tamarix*, *Elaeagnus*) leaf litter. *Freshwater Biology* 53:1012–1020.

Moore GW, Cleverly J, Owens MK. (2008) Nocturnal transpiration in riparian *Tamarix* thickets authenticated by sap flux, eddy covariance and leaf gas exchange measurements. *Tree Physiology* 28:521–528. URL: <http://heronpublishing.com>.

Mortenson SG, Weisberg PJ, Ralston BE. (2008) Do Beavers Promote the Invasion of Non-native *Tamarix* in the Grand Canyon Riparian Zone? *Wetlands* 28(3):666-675.

Nagler PL, Glenn EP, Didan K, Osterberg J, Jordan F, Cunningham J. (2008) Wide-Area Estimates of Stand Structure and Water Use of *Tamarix* spp. on the Lower Colorado River: Implications for Restoration and Water Management Projects. *Restoration Ecology* 16(1):136–145.

Nelson SM, Wydoski R. (2008) Riparian Butterfly (*Papilioidea* and *Hesperioidae*) Assemblages Associated with *Tamarix*-Dominated, Native Vegetation-Dominated, and *Tamarix* Removal Sites along the Arkansas River, Colorado, U.S.A. *Restoration Ecology* 16(1):168–179.

Owens MK, Moore GW. (2007) Saltcedar Water Use: Realistic and Unrealistic Expectations. *Rangeland Ecology and Management* 60(5):553-557.

Schumm SA, Lichty RW. (1963). Erosion and Sedimentation in a Semiarid Environment: Channel widening and flood-plain construction along Cimarron River in southwestern Kansas. 71-88. US Geological Survey professional paper ISSN: 1044-9612.

Shafrroth PB, Briggs MK. (2008) Restoration Ecology and Invasive Riparian Plants: An Introduction to the Special Section on *Tamarix* spp. in Western North America. *Restoration Ecology* 16(1):94–96.

Shafrroth PB, Cleverly JR, Dudley TL, Taylor JP, Van Riper III C, Weeks EP, Stuart JN. (2005) Control of *Tamarix* in the Western United States: Implications for Water Salvage, Wildlife Use, and Riparian Restoration. *Environmental Management* 35(3):231-246.

Shafrroth PB, Beauchamp VB, Briggs MK, Lair K, Scott, ML, Sher AA. (2008) Planning Riparian Restoration in the Context of *Tamarix* Control in Western North America. *Restoration Ecology* 16(1):97–112.

Sher AA, Gieck S, Brown CS, Nissen SJ. (2008) First-Year Responses of Cheatgrass Following *Tamarix* spp. Control and Restoration-Related Disturbances. *Restoration Ecology* 16(1):129–135.

- Sogge MK, Sferra SJ, Paxton EH. (2008) *Tamarix* as Habitat for Birds: Implications for Riparian Restoration in the Southwestern United States. *Restoration Ecology* 16(1):146–154.
- Sorensen MA, Parker DR, Trumble JT. (2009) Effects of pollutant accumulation by the invasive weed saltcedar (*Tamarix ramosissima*) on the biological control agent *Diorhabda elongata* (Coleoptera: Chrysomelidae). *Environmental Pollution* 157:384-391.
- Tallent-Halsell NG, Walker LR. (2002) Responses of *Salix gooddingii* and *Tamarix ramosissima* to Flooding. *Wetlands* 22(4):776-785.
- Van Riper III C, Paxton KL, O'Brien C, Shafrroth PB, McGrath LJ. (2008) Rethinking Avian Response to *Tamarix* on the Lower Colorado River: A Threshold Hypothesis. *Restoration Ecology* 16(1):155–167.
- Walker HA. (2008) Floristics and Physiognomy Determine Migrant Landbird Response to Tamarisk (*Tamarix ramosissima*) Invasion in Riparian Areas. *The Auk* 125(3):520-531.
- Whitcraft CR, Talley DM, Crooks JA, Boland J, Gaskin J. (2007) Invasion of Tamarisk (*Tamarix* spp.) in a Southern California Salt Marsh. *Biological Invasions* 9:875-879.
- Wiesenborn WD. (2005) Biomass of Arthropod Trophic Levels on *Tamarix ramosissima* (Tamaricaceae) Branches. *Environmental Entomology* 34(3):656-663.
- Wiesenborn WD, Heydon SL. (2007) Diets of Breeding Southwestern Willow Flycatchers in Different Habitats. *The Wilson Journal of Ornithology* 119(4):547-557.
- Wiesenborn WD, Heydon SL, Lorenzen K. (2008) Pollen Loads on Adult Insects from Tamarisk Flowers and Inferences about Larval Habitats at Topock Marsh, Arizona. *Journal of the Kansas Entomological Society* 81(1):50-60.
- Zavaleta E. (2000) The Economic Value of Controlling an Invasive Shrub. *Ambio* 29(8):462-467. URL: <http://www.ambio.kva.se>.